

BAILEY TYPE TRANSFORMS AND APPLICATIONS

C.M. Joshi and Yashoverdhan Vyas*

Abstract

The aim of this paper is to establish new series transforms of Bailey type and to show that these Bailey type transforms work as efficiently as the classical one and give not only new q -hypergeometric identities, converting double or triple series into a very-well-poised $_{10}\Phi_9$ or $_{12}\Phi_{11}$ series but can also be utilized to derive new double and triple series Rogers-Ramanujan type identities and corresponding infinite families of Rogers-Ramanujan type identities.

1 Introduction

It is well-known that W. N. Bailey [23, 24] established a series transform known as Bailey's transform and gave a mechanism to derive ordinary and q -hypergeometric identities and Rogers-Ramanujan type identities. Using Bailey's transform, Slater [61, 62] derived 130 Rogers-Ramanujan type identities. It was Andrews [3, 4, 10] and [9], who exploited the Bailey's transform in the form of Bailey pair and Bailey chains to show that all of the 130 identities given by Slater [61, 62, 63] can be embedded in infinite families of multiple series Rogers-Ramanujan type identities.

After Bailey [23, 24] and notably after Andrews [9], a large number of Mathematicians have worked on Bailey's transform in the form of Bailey lemma and Bailey chain to make applications in the theory of generalized hypergeometric series, number theory, partition theory, combinatorics, physics and computer-algebra (see [1, 2, 5, 6, 7, 8, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 25, 26, 28, 29, 31, 32, 33, 34, 35, 36, 38, 41, 42, 43, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 64, 66, 67, 68, 74, 73, 70, 69, 72] and [71]).

A survey of all the above mentioned papers reveals that the work done by Bailey [23, 24] and Andrews [9] may be considered as basic and fundamental in this field. At this point there arises one natural question,

*He is grateful to the **Council of Scientific and Industrial Research (CSIR), New Delhi, INDIA** for providing the Senior Research Fellowship.

“Can there be other series transforms of Bailey type, which not only produce new ordinary and q -hypergeometric identities and Rogers-Ramanujan type identities but can also be iterated to provide multi sum versions of the Rogers-Ramanujan type identities?”

This paper provides the answer in affirmative.

In this paper, first we establish two new series transform of Bailey type in Section 2 as theorems 2.1 and 2.2 and then using these theorems we derive five new q -hypergeometric identities converting double or triple q -hypergeometric series into a very-well-poised $_{10}\Phi_9$ or $_{12}\Phi_{11}$ series in Section 3. In Section 4, we establish the first Bailey type lemma and using it derive four new double series Rogers-Ramanujan type identities of modulo 7 and 5 and embed these in multiple series Rogers-Ramanujan type identities of modulo $4k + 3$ and $4k + 1$. In Section 5, we establish second Bailey type lemma and then using it derive two new triple series Rogers-Ramanujan type identities of modulo 9 and embed these in multiple series Rogers-Ramanujan type identities of modulo $6s + 3$. During the presentation of both lemmas in Section 4 and 5, we establish only, the main results corresponding to Andrews [9, 10] presentation of Bailey lemma.

It may be noted that the double and triple series Rogers-Ramanujan type identities presented in this paper do not arise as merely “one or two level up” in the standard Bailey chain from well-known single sum identities. Furthermore, the original Rogers-Ramanujan identities (see [39, 45, 46]) were also of modulus 5 but were single sum identities. A modulo 7 identity with single sum is by Rogers [47, Eq. 6; p. 331] and as a one level up is by Andrews [4, Eq. (1.8); p. 4083]. Andrews [4] obtained his identity of modulo 7 by particularizing the general multi sum odd moduli identities to double series case. After comparing our identities with the results of references [34, 35, 36, 55, 56, 57, 58, 66, 67] and so forth, we can say that the identities derived in this paper are new and most attractive and symmetric among all the double and triple series Rogers-Ramanujan type identities derived previously.

The main tools in developing these transforms are two series rearrangements (2.5) and (2.10). Some of the instances of use of (2.5) and (2.10) may be found in the works of Burchnell and Chaundy [27], Shankar [48] and Srivastava and Manocha [65, p. 335]. The notation for double q -Kampé de Fériet in the eqn (3.1) and (3.3) is from [65, Eq. (282), (283); p. 349] and of a triple series $\Phi^{(3)}$ in eqn (3.5) is simply a q -analogue of a particular case of the Srivastava’s general triple series [65, Eq. (14), (15); p. 44]. In view of notational difficulties mentioned in [65, pp. 270-274], the double non q -Kampé de Fériet series in eqns (3.2) and (3.4) have been written explicitly without using any notation. However, notation of q -analogue of Srivastava-Daoust’s series [65, Eq. 284; p. 350] may be used in these cases. Remaining definitions and notations are from [33, 10] .

2 Bailey type transforms

2.1 First Bailey type transform

Theorem 2.1

$$\text{If } \beta_{(n,l)} = \sum_{m=0}^{\min.(n,l)} \alpha_m u_{n-m} u'_{l-m} v_{n+m} v'_{l+m} t_{n-l} w_{l+n} \quad (2.1)$$

$$\text{and } \gamma_m = \sum_{n=m}^{\infty} \sum_{l=m}^{\infty} \delta_n \delta'_l u_{n-m} u'_{l-m} v_{n+m} v'_{l+m} t_{n-l} w_{l+n} \quad (2.2)$$

then, subject to convergence conditions

$$\sum_{m=0}^{\infty} \alpha_m \gamma_m = \sum_{n,l=0}^{\infty} \beta_{(n,l)} \delta_n \delta'_l, \quad (2.3)$$

where, $\alpha_r, \delta_r, u_r, v_r, \delta'_r, u'_r, v'_r, t_r$ and w_r are any functions of r only.

Proof :

Observe that

$$\sum_{m=0}^{\infty} \alpha_m \gamma_m = \sum_{m=0}^{\infty} \sum_{n=m}^{\infty} \sum_{l=m}^{\infty} \alpha_m \delta_n \delta'_l u_{n-m} u'_{l-m} v_{n+m} v'_{l+m} t_{n-l} w_{l+n}. \quad (2.4)$$

If this double series is convergent, then using [30; p. 10, lemma 3], viz.,

$$\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} A(m, n, l) = \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{m=0}^{\min.(n,l)} A(m, n-m, l-m), \quad (2.5)$$

in (2.4) after the replacement of n by $n+m$ and of l by $l+m$, we get

$$\begin{aligned} \sum_{m=0}^{\infty} \alpha_m \gamma_m &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{m=0}^{\min.(n,l)} \alpha_m \delta_n \delta'_l u_{n-m} u'_{l-m} v_{n+m} v'_{l+m} t_{n-l} w_{l+n} \\ &= \sum_{n,l=0}^{\infty} \beta_{(n,l)} \delta_n \delta'_l. \end{aligned}$$

2.2 Second Bailey type transform

Theorem 2.2

$$\text{If } \beta_{(n,l,k)} = \sum_{m=0}^{\min.(n,l,k)} \alpha_m u_{n-m} u'_{l-m} u''_{k-m} v_{n+m} v'_{l+m} v''_{k+m} \quad (2.6)$$

$$\text{and } \gamma_m = \sum_{n=m}^{\infty} \sum_{l=m}^{\infty} \sum_{k=m}^{\infty} \delta_n \delta'_l \delta''_k u_{n-m} u'_{l-m} u''_{k-m} v_{n+m} v'_{l+m} v''_{k+m} \quad (2.7)$$

then, subject to convergence conditions

$$\sum_{m=0}^{\infty} \alpha_m \gamma_m = \sum_{n,l,k=0}^{\infty} \beta_{(n,l,k)} \delta_n \delta'_l \delta''_k \quad (2.8)$$

where, $\alpha_r, \delta_r, u_r, v_r, \delta'_r, u'_r, v'_r, \delta''_r, u''_r, v''_r$ are any functions of r only.

Proof :

Observe that

$$\begin{aligned} \sum_{m=0}^{\infty} \alpha_m \gamma_m &= \\ \sum_{m=0}^{\infty} \sum_{n=m}^{\infty} \sum_{l=m}^{\infty} \sum_{k=m}^{\infty} \alpha_m \delta_n \delta'_l \delta''_k u_{n-m} u'_{l-m} u''_{k-m} v_{n+m} v'_{l+m} v''_{k+m}. \end{aligned} \quad (2.9)$$

If this double series is convergent, then using [30; p. 10, lemma 3], viz.,

$$\sum_{m=0}^{\infty} \sum_{n,l,k=0}^{\infty} A(m, n, l, k) = \sum_{n,l,k=0}^{\infty} \sum_{m=0}^{\min.(n,l,k)} A(m, n-m, l-m, k-m), \quad (2.10)$$

in (2.8) after the replacement of n, l and k by $n+m, l+m$ and $k+m$ respectively, we get

$$\begin{aligned} \sum_{m=0}^{\infty} \alpha_m \gamma_m &= \\ \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} \sum_{m=0}^{\min.(n,l,k)} \alpha_m \delta_n \delta'_l \delta''_k u_{n-m} u'_{l-m} u''_{k-m} v_{n+m} v'_{l+m} v''_{k+m} \\ &= \sum_{n,l,k=0}^{\infty} \beta_{(n,l,k)} \delta_n \delta'_l \delta''_k. \end{aligned}$$

Here, it may be noted that in these two theorems various new sequences of other combinations of involved summation indices may be introduced in their $\beta_{(n,l)}$ or $\beta_{(n,l,k)}$ and γ_m . But we have presented these theorems with only those sequences, which have been used at least once in this paper. Further, it may be possible in many cases to decide the new expressions for $\alpha_r, \delta_r, u_r, v_r, \delta'_r, u'_r, v'_r, \delta''_r, u''_r, v''_r, t_r$ and w_r , which yield closed forms for $\beta_{(n,l)}$ or $\beta_{(n,l,k)}$ and γ_m , using one or more of the known summation theorems. Thus, many more new results may be discovered. But we have discussed certain cases which use the well-known classical summation theorems [33, Appendix, II. 5, 6, 12, 21, 22] only. It may be emphasized that there may be other various appropriate cases to have the closed forms for $\beta_{(n,l)}$ or $\beta_{(n,l,k)}$ and γ_m in theorems 2.1 and 2.2.

Further, many series rearrangements, similar to (2.5) and (2.10), may also be utilized to discuss other Bailey type transforms and their applications.

3 New q -hypergeometric identities

In the Bailey type transforms, discussed in the previous Section as theorems 2.1 and 2.2, we observed five expressions for $\alpha_r, \delta_r, u_r, v_r, \delta'_r, u'_r, v'_r, \delta''_r, u''_r, v''_r, t_r$ and w_r , which yield closed forms for $\beta_{(n,l)}$ or $\beta_{(n,l,k)}$ and γ_m and lead to five new q -hypergeometric identities. The identities are as follows:

$$\begin{aligned} & \Phi_{1:2;2}^{1:3;2} \left[\begin{array}{c} \frac{qa}{d} : \quad b, c, q^{-M}; \quad b', c', q^{-N} \\ qa : \quad \frac{qa}{d}, \frac{bcq^{-M}}{a}; \quad \frac{qa}{d}, \frac{b'c'q^{-N}}{a} \end{array} ; q, q \right] \\ &= \frac{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_M \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_N}{\left(qa, \frac{qa}{bc}; q\right)_M \left(qa, \frac{qa}{b'c'}; q\right)_N} \times {}_{10}W_9 \left(a; d, b, c, b', c', q^{-M}, q^{-N}; q, \frac{a^3 q^{3+M+N}}{b c b' c' d} \right), \end{aligned} \quad (3.1)$$

$$\begin{aligned} & \sum_{n,l=0}^{\infty} \frac{\left(\frac{A}{a}; q\right)_{n-l} \left(\frac{AB}{a}; q\right)_{n+l}}{\left(\frac{qa}{B}; q\right)_{n-l} (qa; q)_{n+l}} \frac{\left(A, q\sqrt{A}, -q\sqrt{A}, b, c, \frac{Aaq^{1+M}}{b c}, \frac{qa}{B}, q^{-M}; q\right)_n}{\left(\sqrt{A}, -\sqrt{A}, \frac{qA}{b}, \frac{qA}{c}, \frac{b c q^{-M}}{a}, \frac{AB}{a}, A q^{1+M}; q\right)_n} \\ & \times \frac{\left(B, q\sqrt{B}, -q\sqrt{B}, b', c', \frac{Baq^{1+N}}{b' c'}, \frac{qa}{A}, q^{-N}; q\right)_l q^n \left(\frac{qAB}{a^2}\right)^l}{\left(\sqrt{B}, -\sqrt{B}, \frac{qB}{b'}, \frac{qB}{c'}, \frac{b' c' q^{-N}}{a}, \frac{AB}{a}, B q^{1+N}; q\right)_l (q; q)_n (q; q)_l} \\ &= \frac{\left(qA, \frac{qa}{b}, \frac{qa}{c}, \frac{qA}{bc}; q\right)_M}{\left(qa, \frac{qA}{b}, \frac{qA}{c}, \frac{qa}{bc}; q\right)_M} \cdot \frac{\left(qB, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qB}{b'c'}; q\right)_N}{\left(qa, \frac{qB}{b'}, \frac{qB}{c'}, \frac{qa}{b'c'}; q\right)_N} \\ & \times {}_{12}W_{11} \left(a; \frac{qa^2}{AB}, b, c, b', c', \frac{Aaq^{1+M}}{b c}, \frac{Baq^{1+N}}{b' c'}, q^{-M}, q^{-N}; q, q \right), \end{aligned} \quad (3.2)$$

$$\Phi_{2:1;7}^{2:2;8} \left[\begin{array}{c} b', \frac{qa}{d} : \quad c', q^{-M}; \quad \sqrt{qb'}, -\sqrt{qb'}, b, c, \frac{ab' q^{1+N}}{b c}, \frac{b' c'}{a}, \frac{b' q^{-M}}{a}, q^{-N} \\ qa, \frac{b' c' q^{-M}}{a} : \quad \frac{qa}{d}; \quad \sqrt{b'}, -\sqrt{b'}, \frac{qb'}{b}, \frac{qb'}{c}, \frac{bcq^{-N}}{a}, b' q^{1+N}, \frac{qa}{d} \end{array} ; q, q \right]$$

$$\begin{aligned}
&= \frac{\left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_M}{\left(qa, \frac{qa}{b' c'}; q\right)_M} \cdot \frac{\left(qb', \frac{qa}{b}, \frac{qa}{c}, \frac{qb'}{bc}; q\right)_N}{\left(qa, \frac{qb'}{b}, \frac{qb'}{c}, \frac{qa}{bc}; q\right)_N} \\
&\times {}_{10}W_9 \left(a; d, b, c, c', \frac{q^{1+N}ab'}{bc}, q^{-M}, q^{-N}; q, \frac{q^{2+M}a^2}{b' c'}\right) \quad (3.3)
\end{aligned}$$

$$\begin{aligned}
&\sum_{n,l=0}^{\infty} \frac{\left(\frac{b'}{A}; q\right)_{n-l} (b'; q)_{n+l}}{\left(\frac{qa}{A}; q\right)_{n-l} (qa; q)_{n+l}} \\
&\cdot \frac{\left(q\sqrt{b'}, -q\sqrt{b'}, b, c, \frac{b' q^{-M}}{a}, \frac{b' c'}{a}, \frac{b' a q^{1+N}}{b c}, \frac{b' c' q^{-M}}{A}, \frac{qa}{A}, q^{-N}; q\right)_n}{\left(\sqrt{b'}, -\sqrt{b'}, \frac{b' q^{-M}}{A}, \frac{b' c'}{A}, \frac{qb'}{b}, \frac{qb'}{c}, \frac{b' c' q^{-M}}{a}, \frac{bcq^{-N}}{a}, b' q^{1+N}; q\right)_n} \\
&\cdot \frac{\left(q\sqrt{A}, -q\sqrt{A}, c', A, \frac{Aa q^{1+M}}{b' c'}, q^{-M}; q\right)_l}{\left(\sqrt{A}, -\sqrt{A}, \frac{qA}{c'}, \frac{b' c' q^{-M}}{a}, Aq^{1+M}; q\right)_l} \cdot \frac{q^n \left(\frac{q^{2+M} a^2}{b' c'}\right)^l}{(q; q)_n (q; q)_l} \\
&= \frac{\left(qA, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qA}{b' c'}; q\right)_M}{\left(qa, \frac{qA}{b'}, \frac{qA}{c'}, \frac{qa}{b' c'}; q\right)_M} \cdot \frac{\left(qb', \frac{qa}{b}, \frac{qa}{c}, \frac{qb'}{bc}; q\right)_N}{\left(qa, \frac{qb'}{b}, \frac{qb'}{c}, \frac{qa}{bc}; q\right)_N} \\
&\times {}_{10}W_9 \left(a; b, c, c', \frac{a^2 q^{1+M}}{b' c'}, \frac{ab' q^{1+N}}{bc}, q^{-M}, q^{-N}; q, \frac{qc'}{aq^M}\right) \quad (3.4)
\end{aligned}$$

and

$$\begin{aligned}
&\Phi^{(3)} \left[\begin{array}{ccccccc} qa :: & : & : & : & b, c, q^{-N}; & b', c', q^{-L}; & b'', c'', q^{-K} \\ & :: & qa : & qa : & qa : & \frac{bcq^{-N}}{a}; & \frac{b' c' q^{-L}}{a}; & \frac{b'' c'' q^{-K}}{a} & ; q; q, q, q \end{array} \right] \\
&= \frac{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_N}{\left(qa, \frac{qa}{bc}; q\right)_N} \frac{\left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_L}{\left(qa, \frac{qa}{b' c'}; q\right)_L} \frac{\left(\frac{qa}{b''}, \frac{qa}{c''}; q\right)_K}{\left(qa, \frac{qa}{b'' c''}; q\right)_K} \\
&\times {}_{12}W_{11} \left(a; b, c, b', c', b'', c'', q^{-N}, q^{-L}, q^{-K}; q, \frac{a^4 q^{N+L+K+4}}{b c b' c' b'' c''}\right). \quad (3.5)
\end{aligned}$$

Proof of (3.1)

For the choice

$$\alpha_r = \frac{(a, q\sqrt{a}, -q\sqrt{a}, d; q)_r q^{r^2-r} \left(\frac{aq^3}{d}\right)^r}{\left(\sqrt{a}, -\sqrt{a}, \frac{qa}{d}, q; q\right)_r}, \delta_r = \frac{(b, c, q^{-M}; q)_r}{\left(\frac{bcq^{-M}}{a}; q\right)_r},$$

$$\delta'_r = \frac{(b', c', q^{-N}; q)_r}{\left(\frac{b' c' q^{-N}}{a}; q\right)_r}, v_r = v'_r = \frac{1}{(qa; q)_r}, u_r = u'_r = \frac{q^r}{(q; q)_r},$$

and $t_r = w_r = 1$ in (2.1) and (2.2), the q -Pfaff-Saalschütz sum [33, p. 237, (II. 12)] and the terminating very-well-poised ${}_6\Phi_5$ sum [33, p. 238, (II. 21)] can be used to have

$$\beta_{(n,l)} = \frac{\left(\frac{qa}{d}; q\right)_{n+l} q^{n+l}}{(qa; q)_{n+l} \left(\frac{qa}{d}, q; q\right)_n \left(\frac{qa}{d}, q; q\right)_l}$$

and

$$\gamma_m = \frac{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_M \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_N (b, c, b', c', q^{-M}, q^{-N}; q)_m q^{-m^2+m}}{\left(qa, \frac{qa}{bc}; q\right)_M \left(qa, \frac{qa}{b'c'}; q\right)_N \left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, aq^{1+M}, aq^{1+N}; q\right)_m} \left(\frac{a^2 q^{M+N}}{b c b' c'}\right)^m.$$

Substituting these values in (2.3), we obtain the result (3.1).

Proof of (3.2)

For the choice

$$\alpha_r = \frac{\left(a, q\sqrt{a}, -q\sqrt{a}, \frac{qa^2}{AB}; q\right)_r \left(\frac{qAB}{a^2}\right)^r}{\left(\sqrt{a}, -\sqrt{a}, \frac{AB}{a}, q; q\right)_r},$$

$$\delta_r = \frac{\left(q\sqrt{A}, -q\sqrt{A}, b, c, \frac{Aaq^{1+M}}{bc}, q^{-M}; q\right)_r}{\left(\sqrt{A}, -\sqrt{A}, \frac{qA}{b}, \frac{qA}{c}, \frac{bcq^{-M}}{a}, Aq^{1+M}; q\right)_r},$$

$$\delta'_r = \frac{\left(q\sqrt{B}, -q\sqrt{B}, b', c', \frac{Baq^{1+N}}{b'c'}, q^{-N}; q \right)_r}{\left(\sqrt{B}, -\sqrt{B}, \frac{qB}{b'}, \frac{qB}{c'}, \frac{b'c'q^{-N}}{a}, Bq^{1+N}; q \right)_r},$$

$$v_r = \frac{(A; q)_r}{(qa; q)_r}, v'_r = \frac{(B; q)_r}{(qa; q)_r}, u_r = \frac{\left(\frac{A}{a}; q \right)_r q^r}{(q; q)_r}, u'_r = \frac{\left(\frac{B}{a}; q \right)_r q^r}{(q; q)_r},$$

and $t_r = w_r = 1$ in (2.1) and (2.2) and using the Jackson's q -analogue of Dougall's ${}_7F_6$ sum [33, p. 238, (II. 22)] we can obtain

$$\beta_{(n,l)} = \frac{\left(\frac{A}{a}; q \right)_{n-l} \left(\frac{AB}{a}; q \right)_{n+l} \left(A, \frac{qa}{B}; q \right)_n \left(B, \frac{qa}{A}; q \right)_l}{\left(\frac{qa}{B}; q \right)_{n-l} (qa; q)_{n+l} \left(\frac{AB}{a}, q; q \right)_n \left(\frac{AB}{a}, q; q \right)_l} q^n \left(\frac{AB}{a^2} \right)^l$$

and

$$\gamma_m = \frac{\left(qA, \frac{qa}{b}, \frac{qa}{c}, \frac{qA}{bc}; q \right)_M \left(qB, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qB}{b'c'}; q \right)_N}{\left(qa, \frac{qA}{b}, \frac{qA}{c}, \frac{qa}{bc}; q \right)_M \left(qa, \frac{qB}{b'}, \frac{qB}{c'}, \frac{qa}{b'c'}; q \right)_N}$$

$$\times \frac{\left(b, c, \frac{Aaq^{1+M}}{bc}, q^{-M}, b', c', \frac{Baq^{1+N}}{b'c'}, q^{-N}; q \right)_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{bcq^{-M}}{A}, aq^{1+M}, \frac{qa}{b'}, \frac{qa}{c'}, \frac{b'c'q^{-N}}{B}, aq^{1+N}; q \right)_m} \left(\frac{a^2}{AB} \right)^m.$$

Substituting these values in (2.3), we obtain the result (3.2).

Proof of (3.3)

For the choice

$$\alpha_r = \frac{(a, q\sqrt{a}, -q\sqrt{a}, d; q)_r q^{r^2} \left(\frac{q^2 a}{d} \right)^r}{\left(\sqrt{a}, -\sqrt{a}, \frac{qa}{d}, q; q \right)_r},$$

$$\delta_r = \frac{\left(q\sqrt{b'}, -q\sqrt{b'}, b, c, \frac{ab'q^{1+N}}{bc}, q^{-N}, \frac{b'c'}{a}, \frac{b'q^{-M}}{a}; q \right)_r}{\left(\sqrt{b'}, -\sqrt{b'}, \frac{qb'}{b}, \frac{qb'}{c}, \frac{bcq^{-N}}{a}, b'q^{1+N}; q \right)_r},$$

$$v_r = \frac{1}{(qa; q)_r} = v'_r, u_r = \frac{q^r}{(q; q)_r} = u'_r, w_r = \frac{(b'; q)_r}{\left(\frac{b' c' q^{-M}}{a}; q\right)_r},$$

$\delta'_r = (c', q^{-M}; q)_r$ and $t_r = 1$ in (2.1) and (2.2), the terminating very-well-poised ${}_6\Phi_5$ sum [33, p. 238, (II. 21)], the q -Pfaff-Saalschütz sum [33, p. 237, (II. 12)] and the Jackson's q -analogue of Dougall's ${}_7F_6$ sum [33, p. 238, (II. 22)] leads to

$$\beta_{(n,l)} = \frac{\left(b', \frac{qa}{d}; q\right)_{n+l}}{\left(aq, \frac{b' c' q^{-M}}{a}; q\right)_{n+l}} \frac{q^{n+l}}{\left(\frac{qa}{d}, q; q\right)_n \left(\frac{qa}{d}, q; q\right)_l}$$

and

$$\begin{aligned} \gamma_m &= \frac{\left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_M \left(qb', \frac{qa}{b}, \frac{qa}{c}, \frac{qb'}{bc}; q\right)_N}{\left(qa, \frac{qa}{b' c'}; q\right)_M \left(qa, \frac{qb'}{b}, \frac{qb'}{c}, \frac{qa}{bc}; q\right)_N} \\ &\times \frac{\left(b, c, c', \frac{ab' q^{1+N}}{bc}, q^{-M}, q^{-N}; q\right)_m \left(\frac{a}{b' c'}\right)^m q^{-m^2+mM}}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{c'}, \frac{bcq^{-N}}{b'}, aq^{1+M}, aq^{1+N}; q\right)_m} \end{aligned}$$

Substituting these values in (2.3), we obtain the result (3.3).

Proof of (3.4)

For the choice

$$\begin{aligned} \alpha_r &= \frac{\left(a, q\sqrt{a}, -q\sqrt{a}, \frac{a^2 q^{1+M}}{b' c'}; q\right)_r \left(\frac{qb' c'}{a^2 q^M}\right)^r}{\left(\sqrt{a}, -\sqrt{a}, \frac{b' c' q^{-M}}{a}, q; q\right)_r}, \\ \delta_r &= \frac{\left(q\sqrt{b'}, -q\sqrt{b'}, b, c, \frac{ab' q^{1+N}}{bc}, q^{-N}, \frac{b' c'}{a}, \frac{b' q^{-M}}{a}; q\right)_r}{\left(\sqrt{b'}, -\sqrt{b'}, \frac{qb'}{b}, \frac{qb'}{c}, \frac{bcq^{-N}}{a}, b' q^{1+N}; q\right)_r}, \end{aligned}$$

$$\delta'_r = \frac{\left(q\sqrt{A}, -q\sqrt{A}, c', q^{-M}; q\right)_r}{\left(\sqrt{A}, -\sqrt{A}, \frac{qA}{c'}, Aq^{1+M}; q\right)_r}, w_r = \frac{(b'; q)_r}{\left(\frac{b' c' q^{-M}}{a}; q\right)_r}, t_r = \frac{\left(\frac{b'}{A}; q\right)_r}{\left(\frac{b' c' q^{-M}}{Aa}; q\right)_r},$$

$$v_r = \frac{\left(\frac{b' c' q^{-M}}{A}; q\right)_r}{(qa; q)_r}, v'_r = \frac{(A; q)_r}{(qa; q)_r}, u_r = \frac{\left(\frac{b' c' q^{-M}}{Aa}; q\right)_r q^r}{(q; q)_r} \text{ and } u'_r = \frac{\left(\frac{A}{a}; q\right)_r q^r}{(q; q)_r},$$

in (2.1) and (2.2), the Jackson's q -analogue of Dougall's ${}_7F_6$ sum [33, p. 238, (II. 22)] gives rise to

$$\beta_{(n,l)} = \frac{\left(\frac{a}{A}; q\right)_{n-l} (b'; q)_{n+l} \left(\frac{b' c' q^{-M}}{A}, \frac{qa}{A}; q\right)_n \left(A, \frac{aAq^{1+M}}{b' c'}; q\right)_l}{\left(\frac{qa}{A}; q\right)_{n-l} (qa; q)_{n+l} \left(\frac{b' c' q^{-M}}{A}, q; q\right)_n \left(\frac{b' c' q^{-M}}{A}, q; q\right)_l} q^n \left(\frac{b' c' q^{-M}}{a^2}\right)^l$$

and

$$\gamma_m = \frac{\left(qA, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qA}{b' c'}; q\right)_M \left(qb', \frac{qa}{b}, \frac{qa}{c}, \frac{qb'}{bc}; q\right)_N}{\left(qa, \frac{qA}{b'}, \frac{qA}{c'}, \frac{qa}{b' c'}; q\right)_M \left(qa, \frac{qb'}{b}, \frac{qb'}{c}, \frac{qa}{bc}; q\right)_N}$$

$$\times \frac{\left(b, c, c', \frac{b' a q^{1+N}}{bc}, q^{-M}, q^{-N}; q\right)_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{c'}, \frac{bcq^{-N}}{b'}, aq^{1+M}, aq^{1+N}; q\right)_m} \left(\frac{a}{b'}\right)^m.$$

Substituting these values in (2.3), we obtain the result (3.4).

Proof of (3.5)

For the choice

$$\alpha_r = \frac{(a, q\sqrt{a}, -q\sqrt{a}; q)_r a^r q^{4r} q^{3\binom{r}{2}}}{(\sqrt{a}, -\sqrt{a}, q; q)_r}, \delta_r = \frac{(b, c, q^{-N}; q)_r}{\left(\frac{bcq^{-N}}{a}; q\right)_r},$$

$$\delta'_r = \frac{(b', c', q^{-L}; q)_r}{\left(\frac{b' c' q^{-L}}{a}; q\right)_r}, \quad \delta''_r = \frac{(b'', c'', q^{-K}; q)_r}{\left(\frac{b'' c'' q^{-K}}{a}; q\right)_r},$$

$$v_r = v'_r = v''_r = \frac{1}{(qa; q)_r}, \quad \text{and} \quad u_r = u'_r = u''_r = \frac{q^r}{(q; q)_r},$$

in (2.6) and (2.7), the q -Pfaff-Saalschütz sum [33, p. 237, (II. 12)] and the terminating very-well-poised ${}_6\Phi_5$ sum [33, p. 238, (II. 21)] readily yields

$$\beta_{(n,l,k)} = \frac{(qa; q)_{n+l+k} q^{n+l+k}}{(qa; q)_{n+l} (qa; q)_{n+k} (qa; q)_{l+k} (q; q)_n (q; q)_l (q; q)_k}$$

and

$$\gamma_m = \frac{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_N \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_L \left(\frac{qa}{b''}, \frac{qa}{c''}; q\right)_K}{\left(qa, \frac{qa}{bc}; q\right)_N \left(qa, \frac{qa}{b'c'}; q\right)_L \left(qa, \frac{qa}{b''c''}; q\right)_K}$$

$$\frac{(b, c, b', c', b'', c'', q^{-N}, q^{-L}, q^{-K}; q)_m (-1)^m q^{-3\binom{m}{2}}}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qa}{b''}, \frac{qa}{c''}, aq^{1+N}, aq^{1+L}, aq^{1+K}; q\right)_m} \left(\frac{a^3 q^{N+L+K}}{b c b' c' b'' c''}\right)^m.$$

Substituting these values in (2.8), we obtain the result (3.5).

4 First Bailey type lemma and applications to Rogers-Ramanujan type identities

4.1 First Bailey Type Lemma or FBTL

Theorem 4.1

If for $n, l \geq 0$

$$\beta_{(n,l)} = \sum_{m=0}^{\min.(n,l)} \frac{\alpha_m}{(q; q)_{n-m} (q; q)_{l-m} (aq; q)_{n+m} (aq; q)_{l+m}}, \quad (4.1)$$

then

$$\beta'_{(n,l)} = \sum_{m=0}^{\min.(n,l)} \frac{\alpha'_m}{(q; q)_{n-m} (q; q)_{l-m} (aq; q)_{n+m} (aq; q)_{l+m}}, \quad (4.2)$$

where

$$\alpha'_m = \frac{(b, c, b', c'; q)_m \left(\frac{a^2 q^2}{b c b' c'}\right)^m \alpha_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}; q\right)_m} \quad (4.3)$$

and

$$\beta'_{(M,N)} = \sum_{n,l=0}^{\infty} \frac{(b, c; q)_n \left(\frac{qa}{bc}; q\right)_{M-n} \left(\frac{qa}{bc}\right)^n (b', c'; q)_l \left(\frac{qa}{b'c'}; q\right)_{N-l} \left(\frac{qa}{b'c'}\right)^l \beta_{(n,l)}}{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_M (q; q)_{M-n} \left(\frac{qa}{b}, \frac{qa}{c}; q\right)_N (q; q)_{N-l}}. \quad (4.4)$$

REMARK: A pair of sequences $(\alpha_m, \beta_{(n,l)})$ related by (4.1) may be called as a “first Bailey type pair” or “FBTP” and the Theorem 4.1 may be rephrased as: If $(\alpha_m, \beta_{(n,l)})$ is a FBTP, so is $(\alpha'_m, \beta'_{(n,l)})$ where this new FBTP is given by (4.3) and (4.4).

Proof:

To prove FBTL, we apply Bailey type transform (2.3) with the choice:

$$\delta_r = \frac{(b, c, q^{-M}; q)_r q^r}{\left(\frac{bcq^{-M}}{a}; q\right)_r}, \delta'_r = \frac{(b', c', q^{-N}; q)_r q^r}{\left(\frac{b'c'q^{-N}}{a}; q\right)_r},$$

$$v_r = v'_r = \frac{1}{(qa; q)_r}, u_r = u'_r = \frac{1}{(q; q)_r},$$

and $t_r = w_r = 1$ then as in the proof of (3.1), we get

$$\gamma_m = \frac{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_M \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_N}{\left(qa, \frac{qa}{bc}; q\right)_M \left(qa, \frac{qa}{b'c'}; q\right)_N} \times \frac{(b, c, b', c', q^{-M}, q^{-N}; q)_m q^{-m^2+m}}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, aq^{1+M}, aq^{1+N}; q\right)_m} \left(\frac{a^2 q^{2+M+N}}{b c b' c'}\right)^m.$$

Now we can prove (4.2) as shown below:

$$\begin{aligned}
& \sum_{m=0}^{\min.(M,N)} \frac{\alpha'_m}{(q; q)_{M-m} (q; q)_{N-m} (aq; q)_{M+m} (aq; q)_{N+m}} \\
&= \sum_{m=0}^{\min.(M,N)} \frac{(b, c, b', c'; q)_m \left(\frac{a^2 q^2}{b c b' c'} \right)^m \alpha_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}; q \right)_m (q; q)_{M-m} (q; q)_{N-m} (aq; q)_{M+m} (aq; q)_{N+m}} \\
&= \frac{1}{(qa, q; q)_M} \frac{1}{(qa, q; q)_N} \\
&\times \sum_{m=0}^{\min.(M,N)} \frac{(b, c, b', c', q^{-M}, q^{-N}; q)_m q^{-m^2+m}}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, aq^{1+M}, aq^{1+N}; q \right)_m} \left(\frac{a^2 q^{M+N}}{b c b' c'} \right)^m \alpha_m \\
&= \frac{\left(\frac{qa}{bc}; q \right)_M}{\left(\frac{qa}{b}, \frac{qa}{c}, q; q \right)_M} \frac{\left(\frac{qa}{b' c'}; q \right)_N}{\left(\frac{qa}{b'}, \frac{qa}{c'}, q; q \right)_N} \sum_{m=0}^{\min.(M,N)} \gamma_m \alpha_m \\
&= \frac{\left(\frac{qa}{bc}; q \right)_M}{\left(\frac{qa}{b}, \frac{qa}{c}, q; q \right)_M} \frac{\left(\frac{qa}{b' c'}; q \right)_N}{\left(\frac{qa}{b'}, \frac{qa}{c'}, q; q \right)_N} \sum_{n=0}^M \sum_{l=0}^N \beta_{(n,l)} \delta_n \delta'_l \quad (\text{by Theorem 4.1})
\end{aligned}$$

$$\begin{aligned}
&= \frac{\left(\frac{qa}{bc}; q\right)_M \left(\frac{qa}{b'c'}; q\right)_N}{\left(\frac{qa}{b}, \frac{qa}{c}, q; q\right)_M \left(\frac{qa}{b'}, \frac{qa}{c'}, q; q\right)_N} \\
&\times \sum_{n=0}^M \sum_{l=0}^N \frac{(b, c, q^{-M}; q)_n q^n (b', c', q^{-N}; q)_l q^l \beta_{(n,l)}}{\left(\frac{bcq^{-M}}{a}; q\right)_n \left(\frac{b'c'q^{-N}}{a}; q\right)_l} \\
&= \beta'_{(M,N)}.
\end{aligned}$$

The last line follows by manipulating q -shifted factorials, which reduces the previous expression to (4.4). $\square$

Now to derive Rogers-Ramanujan type identities, we need the following result obtained by substituting the values of α'_m and $\beta'_{(n,l)}$ from (4.3) and (4.4) into (4.2), viz.

$$\begin{aligned}
&\sum_{n,l=0}^{\infty} \frac{(b, c; q)_n \left(\frac{qa}{bc}; q\right)_{M-n} \left(\frac{qa}{bc}\right)^n (b', c'; q)_l \left(\frac{qa}{b'c'}; q\right)_{N-l} \left(\frac{qa}{b'c'}\right)^l \beta_{(n,l)}}{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_M (q; q)_{M-n} \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_N (q; q)_{N-l}} \\
&= \sum_{m=0}^{\min(M,N)} \frac{(b, c, b', c'; q)_m \left(\frac{a^2 q^2}{bcb'c'}\right)^m \alpha_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}; q\right)_m (q; q)_{M-m} (q; q)_{N-m} (aq; q)_{M+m} (aq; q)_{N+m}} \quad (4.5)
\end{aligned}$$

Now taking $b, c, b', c', M, N \rightarrow \infty$ in (4.5), we get

$$\sum_{n,l=0}^{\infty} a^{n+l} q^{n^2+l^2} \beta_{(n,l)} = \frac{1}{(aq; q)_{\infty}^2} \sum_{m=0}^{\infty} a^{2m} q^{2m^2} \alpha_m, \quad (4.6)$$

for any FBTP $(\alpha_m, \beta_{(n,l)})$. Now, we shall make use of the following two “FBTPs” deduced by us :

$$\begin{aligned}
\alpha_m &= \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m a^m q^{\frac{1}{2}(3m^2-m)}, \\
\beta_{(n,l)} &= \frac{1}{(aq; q)_{n+l} (q; q)_n (q; q)_l},
\end{aligned} \quad (4.7)$$

and

$$\alpha_m = \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m q^{\frac{1}{2}(m^2 - m)},$$

$$\beta_{(n,l)} = \frac{q^{nl}}{(qa; q)_{n+l} (q; q)_n (q; q)_l}.$$
(4.8)

The fact that “FBTPs” $(\alpha_m, \beta_{(n,l)})$ given by (4.7) and (4.8) satisfy (4.1) may be verified by substituting them in (4.1) and then appealing to terminating very-well-poised ${}_6\Phi_5$ sum [33, p. 238, (II. 21)] and taking $d \rightarrow \infty$ or $d \rightarrow 0$.

4.2 New double series Rogers-Ramanujan type identities and corresponding infinite families :

To derive Rogers-Ramanujan type identities, we insert the “FBTPs” (4.7) and (4.8) in (4.6) to get (4.9) and (4.10), respectively, as given below:

$$\sum_{n,l=0}^{\infty} \frac{a^{n+l} q^{n^2+l^2}}{(qa; q)_{n+l} (q; q)_n (q; q)_l}$$

$$= \frac{1}{(aq; q)_{\infty}^2} \sum_{m=0}^{\infty} \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m a^{3m} q^{\frac{1}{2}(7m^2 - m)}, \quad (4.9)$$

$$\sum_{n,l=0}^{\infty} \frac{a^{n+l} q^{n^2+l^2+nl}}{(qa; q)_{n+l} (q; q)_n (q; q)_l}$$

$$= \frac{1}{(aq; q)_{\infty}^2} \sum_{m=0}^{\infty} \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m a^{2m} q^{\frac{1}{2}(5m^2 - m)}, \quad (4.10)$$

and then setting $a = 1$ and $a = q$ in (4.9) and (4.10) and using Jacobi triple product identity, we obtain following four new double series Rogers-Ramanujan type identities:

$$\sum_{n,l=0}^{\infty} \frac{q^{n^2+l^2}}{(q; q)_{n+l} (q; q)_n (q; q)_l} = \frac{1}{(q; q)_{\infty}} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 3 \pmod{7}}}^{\infty} (1 - q^m)^{-1}, \quad (4.11)$$

$$\sum_{n,l=0}^{\infty} \frac{q^{n^2+l^2+n+l}}{(q^2; q)_{n+l} (q; q)_n (q; q)_l} = \frac{1}{(q^2; q)_{\infty}} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{7}}}^{\infty} (1 - q^m)^{-1}, \quad (4.12)$$

$$\sum_{n,l=0}^{\infty} \frac{q^{n^2+l^2+nl}}{(q; q)_{n+l} (q; q)_n (q; q)_l} = \frac{1}{(q; q)_{\infty}} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 2 \pmod{5}}}^{\infty} (1 - q^m)^{-1}, \quad (4.13)$$

and

$$\sum_{n,l=0}^{\infty} \frac{q^{n^2+l^2+n+l+nl}}{(q^2; q)_{n+l} (q; q)_n (q; q)_l} = \frac{1}{(q^2; q)_{\infty}} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{5}}}^{\infty} (1 - q^m)^{-1}. \quad (4.14)$$

Further, like classical Bailey lemma, the idea of repeated application of FBTL may be expressed in the concept of the “first Bailey type chain.” Given a FBTP $(\alpha_m, \beta_{(n,l)})$ we can by FBTL produce new FBTP $(\alpha'_m, \beta'_{(n,l)})$ defined by (4.3) and (4.4). From $(\alpha'_m, \beta'_{(n,l)})$ we can create $(\alpha''_m, \beta''_{(n,l)})$ merely by applying FBTL with $(\alpha'_m, \beta'_{(n,l)})$ as initial FBTP. In this way we create a sequence of FBTPs:

$$(\alpha_m, \beta_{(n,l)}) \rightarrow (\alpha'_m, \beta'_{(n,l)}) \rightarrow (\alpha''_m, \beta''_{(n,l)}) \rightarrow (\alpha'''_m, \beta'''_{(n,l)}) \rightarrow \cdots ;$$

and call it “first Bailey type chain.” Using this chain we can establish:

Theorem 4.2

$$\begin{aligned}
& \sum_{m \geq 0} \frac{(b_1; q)_m (c_1; q)_m (b_2; q)_m (c_2; q)_m \cdots (b_k; q)_m (c_k; q)_m}{\left(\frac{aq}{b_1}; q\right)_m \left(\frac{aq}{c_1}; q\right)_m \left(\frac{aq}{b_2}; q\right)_m \left(\frac{aq}{c_2}; q\right)_m \cdots \left(\frac{aq}{b_k}; q\right)_m \left(\frac{aq}{c_k}; q\right)_m} \\
& \times \frac{(b'_1; q)_m (c'_1; q)_m (b'_2; q)_m (c'_2; q)_m \cdots (b'_k; q)_m (c'_k; q)_m}{\left(\frac{aq}{b'_1}; q\right)_m \left(\frac{aq}{c'_1}; q\right)_m \left(\frac{aq}{b'_2}; q\right)_m \left(\frac{aq}{c'_2}; q\right)_m \cdots \left(\frac{aq}{b'_k}; q\right)_m \left(\frac{aq}{c'_k}; q\right)_m} \\
& \times \frac{(q^{-M}; q)_m (q^{-N}; q)_m}{(aq^{1+M}; q)_m (aq^{1+N}; q)_m} \left(\frac{a^{2k} q^{2k+M+N}}{b_1 c_1 \cdots b_k c_k b'_1 c'_1 \cdots b'_k c'_k} \right)^m q^{-m^2+m} \alpha_m \\
& = \frac{(aq; q)_M \left(\frac{aq}{b_k c_k}; q\right)_M (aq; q)_N \left(\frac{aq}{b'_k c'_k}; q\right)_N}{\left(\frac{aq}{b_k}; q\right)_M \left(\frac{aq}{c_k}; q\right)_M \left(\frac{aq}{b'_k}; q\right)_N \left(\frac{aq}{c'_k}; q\right)_N} \\
& \times \sum_{\substack{n_k \geq \cdots \geq n_1 \geq 0 \\ l_k \geq \cdots \geq l_1 \geq 0}} \frac{(b_k; q)_{n_k} (c_k; q)_{n_k} \cdots (b_1; q)_{n_1} (c_1; q)_{n_1}}{(q; q)_{n_k - n_{k-1}} (q; q)_{n_{k-1} - n_{k-2}} \cdots (q; q)_{n_2 - n_1}} \\
& \times \frac{(b'_k; q)_{l_k} (c'_k; q)_{l_k} \cdots (b'_1; q)_{l_1} (c'_1; q)_{l_1}}{(q; q)_{l_k - l_{k-1}} (q; q)_{l_{k-1} - l_{k-2}} \cdots (q; q)_{l_2 - l_1}} \frac{(q^{-M}; q)_{n_k}}{\left(\frac{b_k c_k q^{-M}}{a}; q\right)_{n_k}} \frac{(q^{-N}; q)_{l_k}}{\left(\frac{b'_k c'_k q^{-N}}{a}; q\right)_{l_k}} \\
& \times \frac{\left(\frac{aq}{b_{k-1} c_{k-1}}; q\right)_{n_k - n_{k-1}} \cdots \left(\frac{aq}{b_1 c_1}; q\right)_{n_2 - n_1}}{\left(\frac{aq}{b_{k-1}}; q\right)_{n_k} \left(\frac{aq}{c_{k-1}}; q\right)_{n_k} \cdots \left(\frac{aq}{b_1}; q\right)_{n_2} \left(\frac{aq}{c_1}; q\right)_{n_2}}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{\left(\frac{aq}{b'_{k-1} c'_{k-1}}; q\right)_{l_k - l_{k-1}} \cdots \left(\frac{aq}{b'_1 c'_1}; q\right)_{l_2 - l_1}}{\left(\frac{aq}{b'_{k-1}}; q\right)_{l_k} \left(\frac{aq}{c'_{k-1}}; q\right)_{l_k} \cdots \left(\frac{aq}{b'_1}; q\right)_{l_2} \left(\frac{aq}{c'_1}; q\right)_{l_2}} \\
& \times q^{n_1 + n_2 + \cdots + n_k + l_1 + l_2 + \cdots + l_k} a^{n_1 + n_2 + \cdots + n_{k-1} + l_1 + l_2 + \cdots + l_{k-1}} \\
& \times (b_{k-1} c_{k-1})^{-n_{k-1}} \cdots (b_1 c_1)^{-n_1} (b'_{k-1} c'_{k-1})^{-l_{k-1}} \cdots (b'_1 c'_1)^{-l_1} \beta_{(n_1, l_1)}. \quad (4.15)
\end{aligned}$$

Proof :

Equation (4.15) is the result of a k -fold iteration of FBTL with the choice $b = b_i$, $c = c_i$, $b' = b'_i$, $c' = c'_i$ at the i th step. $\square$

Theorem 4.3

If $(\alpha_m, \beta_{(n,l)})$ is a FBTP (i.e. related by (4.1)), then

$$\begin{aligned}
& \sum_{\substack{n_k \geq \cdots \geq n_1 \geq 0 \\ l_k \geq \cdots \geq l_1 \geq 0}} \frac{a^{n_1 + n_2 + \cdots + n_k + l_1 + l_2 + \cdots + l_k} q^{n_1^2 + n_2^2 + \cdots + n_k^2 + l_1^2 + l_2^2 + \cdots + l_k^2} \beta_{(n_1, l_1)}}{(q; q)_{n_k - n_{k-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_k - l_{k-1}} \cdots (q; q)_{l_2 - l_1}} \\
& = \frac{1}{(aq; q)_\infty^2} \sum_{m=0}^{\infty} q^{2km^2} a^{2km} \alpha_m. \quad (4.16)
\end{aligned}$$

Proof :

Let $M, N, b_1, \dots, b_k, c_1, \dots, c_k, b'_1, \dots, b'_k, c'_1, \dots, c'_k$ all tend to infinity in Theorem 4.2. $\square$

Theorem 4.3 may now be used to embed the double series Rogers-Ramanujan type identities (4.11) - (4.14) in an infinite family of such identities :

$$\begin{aligned}
& \sum_{\substack{n_k \geq \cdots \geq n_1 \geq 0 \\ l_k \geq \cdots \geq l_1 \geq 0}} \frac{q^{n_1^2 + n_2^2 + \cdots + n_k^2 + l_1^2 + l_2^2 + \cdots + l_k^2}}{(q; q)_{n_k - n_{k-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_k - l_{k-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{1}{(q; q)_{n_1 + l_1} (q; q)_{n_1} (q; q)_{l_1}} = \frac{1}{(q; q)_\infty} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm(2k+1) \pmod{4k+3}}}^{\infty} (1 - q^m)^{-1}, \quad (4.17)
\end{aligned}$$

$$\begin{aligned}
& \sum_{\substack{n_k \geq \dots \geq n_1 \geq 0 \\ l_k \geq \dots \geq l_1 \geq 0}} \frac{q^{n_1^2 + n_2^2 + \dots + n_k^2 + l_1^2 + l_2^2 + \dots + l_k^2 + n_1 + n_2 + \dots + n_k + l_1 + l_2 + \dots + l_k}}{(q; q)_{n_k - n_{k-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_k - l_{k-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{1}{(q^2; q)_{n_1 + l_1} (q; q)_{n_1} (q; q)_{l_1}} = \frac{1}{(q^2; q)_\infty} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{4k+3}}}^{\infty} (1 - q^m)^{-1}, \quad (4.18)
\end{aligned}$$

$$\begin{aligned}
& \sum_{\substack{n_k \geq \dots \geq n_1 \geq 0 \\ l_k \geq \dots \geq l_1 \geq 0}} \frac{q^{n_1^2 + n_2^2 + \dots + n_k^2 + l_1^2 + l_2^2 + \dots + l_k^2 + n_1 l_1}}{(q; q)_{n_k - n_{k-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_k - l_{k-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \frac{1}{(q; q)_{n_1 + l_1} (q; q)_{n_1} (q; q)_{l_1}} = \frac{1}{(q; q)_\infty} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 2k \pmod{4k+1}}}^{\infty} (1 - q^m)^{-1} \quad (4.19)
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{\substack{n_k \geq \dots \geq n_1 \geq 0 \\ l_k \geq \dots \geq l_1 \geq 0}} \frac{q^{n_1^2 + n_2^2 + \dots + n_k^2 + l_1^2 + l_2^2 + \dots + l_k^2 + n_1 + n_2 + \dots + n_k + l_1 + l_2 + \dots + l_k + n_1 l_1}}{(q; q)_{n_k - n_{k-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_k - l_{k-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \frac{1}{(q^2; q)_{n_1 + l_1} (q; q)_{n_1} (q; q)_{l_1}} = \frac{1}{(q^2; q)_\infty} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{4k+1}}}^{\infty} (1 - q^m)^{-1}. \quad (4.20)
\end{aligned}$$

Equation (4.17) follows by using FBTP defined by (4.7) in (4.16) and then invoking to Jacobi triple product identity after taking $a = 1$. For (4.18), we follow the same procedure with $a = q$. In the same way we establish (4.19) and (4.20) by using FBTP defined by (4.8).

5 Second Bailey type lemma and applications to Rogers-Ramanujan type identities

5.1 Second Bailey Type Lemma or SBTL

Theorem 5.1

If for $n, l, k \geq 0$

$$\beta_{(n,l,k)} = \sum_{m=0}^{\min.(n,l,k)} \frac{\alpha_m}{(q; q)_{n-m} (q; q)_{l-m} (q; q)_{k-m} (aq; q)_{n+m} (aq; q)_{l+m} (aq; q)_{k+m}}, \quad (5.1)$$

then

$$\beta'_{(n,l,k)} = \sum_{m=0}^{\min.(n,l,k)} \frac{\alpha'_m}{(q; q)_{n-m} (q; q)_{l-m} (q; q)_{k-m} (aq; q)_{n+m} (aq; q)_{l+m} (aq; q)_{k+m}}, \quad (5.2)$$

where

$$\alpha'_m = \frac{(b, c, b', c', b'', c''; q)_m \left(\frac{a^3 q^3}{b c b' c' b'' c''} \right)^m \alpha_m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qa}{b''}, \frac{qa}{c''}; q \right)_m} \quad (5.3)$$

and

$$\begin{aligned} \beta'_{(N,L,K)} = \sum_{n,l,k=0}^{\infty} & \frac{(b, c; q)_n \left(\frac{qa}{bc}; q \right)_{N-n} \left(\frac{qa}{bc} \right)^n (b', c'; q)_l \left(\frac{qa}{b' c'}; q \right)_{L-l} \left(\frac{qa}{b' c'} \right)^l}{\left(\frac{qa}{b}, \frac{qa}{c}; q \right)_N (q; q)_{N-n} \left(\frac{qa}{b'}, \frac{qa}{c'}; q \right)_L (q; q)_{L-l}} \\ & \times \frac{(b'', c''; q)_k \left(\frac{qa}{b'' c''}; q \right)_{K-k} \left(\frac{qa}{b'' c''} \right)^k \beta_{(n,l,k)}}{\left(\frac{qa}{b''}, \frac{qa}{c''}; q \right)_K (q; q)_{K-k}}. \quad (5.4) \end{aligned}$$

REMARK: A pair of sequences $(\alpha_m, \beta_{(n,l,k)})$ related by (5.1) may be called as a “second Bailey type pair” or “SBTP” and the Theorem 5.1 may be rephrased as: If $(\alpha_m, \beta_{(n,l,k)})$ is a SBTP, so is $(\alpha'_m, \beta'_{(n,l,k)})$ where this new SBTP is given by (5.3) and (5.4).

Proof:

To prove SBTL, we apply Bailey type transform (2.8) with the choice:

$$\delta_r = \frac{(b, c, q^{-N}; q)_r q^r}{\left(\frac{bcq^{-N}}{a}; q\right)_r}, \delta'_r = \frac{(b', c', q^{-L}; q)_r q^r}{\left(\frac{b' c' q^{-L}}{a}; q\right)_r}, \delta''_r = \frac{(b'', c'', q^{-K}; q)_r q^r}{\left(\frac{b'' c'' q^{-K}}{a}; q\right)_r},$$

and

$$v_r = v'_r = v''_r = \frac{1}{(qa; q)_r}, u_r = u'_r = u''_r = \frac{1}{(q; q)_r},$$

then following the proof of Theorem 4.1 we can also prove Theorem 5.1 .

Now to derive Rogers-Ramanujan type identities, we need the following result obtained by substituting the values of α'_m and $\beta'_{(n,l,k)}$ from (5.3) and (5.4) into (5.2), viz.,

$$\begin{aligned} & \sum_{n,l,k=0}^{\infty} \frac{(b, c; q)_n \left(\frac{qa}{bc}; q\right)_{N-n} \left(\frac{qa}{bc}\right)^n (b', c'; q)_l \left(\frac{qa}{b' c'}; q\right)_{L-l} \left(\frac{qa}{b' c'}\right)^l}{\left(\frac{qa}{b}, \frac{qa}{c}; q\right)_N (q; q)_{N-n} \left(\frac{qa}{b'}, \frac{qa}{c'}; q\right)_L (q; q)_{L-l}} \\ & \times \frac{(b'', c''; q)_k \left(\frac{qa}{b'' c''}; q\right)_{K-k} \left(\frac{qa}{b'' c''}\right)^k \beta_{(n,l,k)}}{\left(\frac{qa}{b''}, \frac{qa}{c''}; q\right)_K (q; q)_{K-k}} \\ & = \sum_{m=0}^{\min.(N,L,K)} \frac{(b, c, b', c', b'', c''; q)_m \left(\frac{a^3 q^3}{b c b' c' b'' c''}\right)^m}{\left(\frac{qa}{b}, \frac{qa}{c}, \frac{qa}{b'}, \frac{qa}{c'}, \frac{qa}{b''}, \frac{qa}{c''}; q\right)_m} \\ & \times \frac{\alpha_m}{(q; q)_{N-m} (q; q)_{L-m} (q; q)_{K-m} (aq; q)_{N+m} (aq; q)_{L+m} (aq; q)_{K+m}}. \quad (5.5) \end{aligned}$$

Now taking $b, c, b', c', b'', c'', M, N \rightarrow \infty$ in (4.5), we get

$$\sum_{n,l,k=0}^{\infty} a^{n+l+k} q^{n^2+l^2+k^2} \beta_{(n,l,k)} = \frac{1}{(aq; q)_{\infty}^3} \sum_{m=0}^{\infty} a^{3m} q^{3m^2} \alpha_m, \quad (5.6)$$

for any SBTP $(\alpha_m, \beta_{(n,l,k)})$. Now, we shall make use of the following two “SBTP” deduced by us :

$$\alpha_m = \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m a^m q^{\frac{1}{2}(3m^2 - m)}, \quad (5.7)$$

$$\beta_{(n,l,k)} = \frac{(aq; q)_{n+l+k}}{(qa; q)_{n+l} (qa; q)_{n+k} (qa; q)_{l+k} (q; q)_n (q; q)_l (q; q)_k}.$$

The fact that “SBTP” $(\alpha_m, \beta_{(n,l,k)})$ given by (5.7) satisfy (5.1) may be verified by substituting it in (5.1) and then appealing to terminating very-well-poised ${}_6\Phi_5$ sum [33, p. 238, (II. 21)].

5.2 New triple series Rogers-Ramanujan type identities and corresponding infinite families :

To derive Rogers-Ramanujan type identities, we insert the “SBTP” (5.7) in (5.6) to get (5.8) as given below:

$$\begin{aligned} \sum_{n,l,k=0}^{\infty} \frac{a^{n+l+k} q^{n^2+l^2+k^2} (qa; q)_{n+l+k}}{(qa; q)_{n+l} (qa; q)_{n+k} (qa; q)_{l+k} (q; q)_n (q; q)_l (q; q)_k} \\ = \frac{1}{(aq; q)_{\infty}^3} \sum_{m=0}^{\infty} \frac{(a; q)_m (1 - aq^{2m})}{(q; q)_m (1 - a)} (-1)^m a^{4m} q^{\frac{1}{2}(9m^2 - m)} \end{aligned} \quad (5.8)$$

and then setting $a = 1$ and $a = q$ in (5.8) and using Jacobi triple product identity, we obtain following two new triple series Rogers-Ramanujan type identities:

$$\begin{aligned} \sum_{n,l,k=0}^{\infty} \frac{q^{n^2+l^2+k^2} (q; q)_{n+l+k}}{(q; q)_{n+l} (q; q)_{n+k} (q; q)_{l+k} (q; q)_n (q; q)_l (q; q)_l} \\ = \frac{1}{(q; q)_{\infty}^2} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 4 \pmod{9}}}^{\infty} (1 - q^m)^{-1}, \end{aligned} \quad (5.9)$$

$$\begin{aligned} \sum_{n,l,k=0}^{\infty} \frac{q^{n^2+l^2+k^2+n+l+k} (q^2; q)_{n+l+k}}{(q^2; q)_{n+l} (q^2; q)_{n+k} (q^2; q)_{l+k} (q; q)_n (q; q)_l (q; q)_k} \\ = \frac{1}{(q^2; q)_{\infty}^2} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{9}}}^{\infty} (1 - q^m)^{-1}. \end{aligned} \quad (5.10)$$

Further, like classical Bailey lemma and FBTL, the idea of repeated application of SBTL may be expressed in the concept of the “second Bailey type chain” a sequence of SBTPs:

$$(\alpha_m, \beta_{(n,l,k)}) \rightarrow (\alpha'_m, \beta'_{(n,l,k)}) \rightarrow (\alpha''_m, \beta''_{(n,l,k)}) \rightarrow (\alpha'''_m, \beta'''_{(n,l,k)}) \rightarrow \cdots ;$$

Using this chain we can establish:

Theorem 5.2

$$\begin{aligned} & \sum_{m \geq 0} \frac{(b_1; q)_m (c_1; q)_m (b_2; q)_m (c_2; q)_m \cdots (b_s; q)_m (c_s; q)_m}{\left(\frac{aq}{b_1}; q\right)_m \left(\frac{aq}{c_1}; q\right)_m \left(\frac{aq}{b_2}; q\right)_m \left(\frac{aq}{c_2}; q\right)_m \cdots \left(\frac{aq}{b_s}; q\right)_m \left(\frac{aq}{c_s}; q\right)_m} \\ & \times \frac{(b'_1; q)_m (c'_1; q)_m (b'_2; q)_m (c'_2; q)_m \cdots (b'_s; q)_m (c'_s; q)_m}{\left(\frac{aq}{b'_1}; q\right)_m \left(\frac{aq}{c'_1}; q\right)_m \left(\frac{aq}{b'_2}; q\right)_m \left(\frac{aq}{c'_2}; q\right)_m \cdots \left(\frac{aq}{b'_s}; q\right)_m \left(\frac{aq}{c'_s}; q\right)_m} \\ & \times \frac{(b''_1; q)_m (c''_1; q)_m (b''_2; q)_m (c''_2; q)_m \cdots (b''_s; q)_m (c''_s; q)_m}{\left(\frac{aq}{b''_1}; q\right)_m \left(\frac{aq}{c''_1}; q\right)_m \left(\frac{aq}{b''_2}; q\right)_m \left(\frac{aq}{c''_2}; q\right)_m \cdots \left(\frac{aq}{b''_s}; q\right)_m \left(\frac{aq}{c''_s}; q\right)_m} \\ & \times \frac{(q^{-N}; q)_m (q^{-L}; q)_m (q^{-K}; q)_m}{(aq^{1+N}; q)_m (aq^{1+L}; q)_m (aq^{1+K}; q)_m} \\ & \left(\frac{a^{3s} q^{3s+N+L+K}}{b_1 c_1 \cdots b_s c_s b'_1 c'_1 \cdots b'_s c'_s b''_1 c''_1 \cdots b''_s c''_s} \right)^m q^{\frac{3}{2}(-m^2+m)} \alpha_m \\ & = \frac{(aq; q)_N \left(\frac{aq}{b_s c_s}; q\right)_N (aq; q)_L \left(\frac{aq}{b'_s c'_s}; q\right)_L (aq; q)_K \left(\frac{aq}{b''_s c''_s}; q\right)_K}{\left(\frac{aq}{b_s}; q\right)_N \left(\frac{aq}{c_s}; q\right)_N \left(\frac{aq}{b'_s}; q\right)_L \left(\frac{aq}{c'_s}; q\right)_L \left(\frac{aq}{b''_s}; q\right)_K \left(\frac{aq}{c''_s}; q\right)_K} \\ & \times \sum_{\substack{n_s \geq \cdots \geq n_1 \geq 0 \\ l_s \geq \cdots \geq l_1 \geq 0 \\ k_s \geq \cdots \geq k_1 \geq 0}} \frac{(b_s; q)_{n_s} (c_s; q)_{n_s} \cdots (b_1; q)_{n_1} (c_1; q)_{n_1}}{(q; q)_{n_s - n_{s-1}} (q; q)_{n_{s-1} - n_{s-2}} \cdots (q; q)_{n_2 - n_1}} \end{aligned}$$

$$\begin{aligned}
& \times \frac{(b'_s; q)_{l_s} (c'_s; q)_{l_s} \cdots (b'_1; q)_{l_1} (c'_1; q)_{l_1}}{(q; q)_{l_s - l_{s-1}} (q; q)_{l_{s-1} - l_{s-2}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{(b''_s; q)_{k_s} (c''_s; q)_{k_s} \cdots (b''_1; q)_{l_1} (c''_1; q)_{l_1}}{(q; q)_{k_s - k_{s-1}} (q; q)_{k_{s-1} - k_{s-2}} \cdots (q; q)_{k_2 - k_1}} \\
& \times \frac{(q^{-N}; q)_{n_s} (q^{-L}; q)_{l_s} (q^{-K}; q)_{k_s}}{\left(\frac{b_s c_s q^{-N}}{a}; q \right)_{n_s} \left(\frac{b'_s c'_s q^{-L}}{a}; q \right)_{l_s} \left(\frac{b''_s c''_s q^{-K}}{a}; q \right)_{k_s}} \\
& \times \frac{\left(\frac{aq}{b_{s-1} c_{s-1}}; q \right)_{n_s - n_{s-1}} \cdots \left(\frac{aq}{b_1 c_1}; q \right)_{n_2 - n_1}}{\left(\frac{aq}{b_{s-1}}; q \right)_{n_s} \left(\frac{aq}{c_{s-1}}; q \right)_{n_s} \cdots \left(\frac{aq}{b_1}; q \right)_{n_2} \left(\frac{aq}{c_1}; q \right)_{n_2}} \\
& \times \frac{\left(\frac{aq}{b'_{s-1} c'_{s-1}}; q \right)_{l_s - l_{s-1}} \cdots \left(\frac{aq}{b'_1 c'_1}; q \right)_{l_2 - l_1}}{\left(\frac{aq}{b'_{s-1}}; q \right)_{l_s} \left(\frac{aq}{c'_{s-1}}; q \right)_{l_s} \cdots \left(\frac{aq}{b'_1}; q \right)_{l_2} \left(\frac{aq}{c'_1}; q \right)_{l_2}} \\
& \times \frac{\left(\frac{aq}{b''_{s-1} c''_{s-1}}; q \right)_{k_s - k_{s-1}} \cdots \left(\frac{aq}{b''_1 c''_1}; q \right)_{k_2 - k_1}}{\left(\frac{aq}{b''_{s-1}}; q \right)_{k_s} \left(\frac{aq}{c''_{s-1}}; q \right)_{k_s} \cdots \left(\frac{aq}{b''_1}; q \right)_{k_2} \left(\frac{aq}{c''_1}; q \right)_{k_2}} \\
& \times q^{n_1 + n_2 + \cdots + n_s + l_1 + l_2 + \cdots + l_s + k_1 + k_2 + \cdots + k_s} \\
& \times a^{n_1 + n_2 + \cdots + n_{s-1} + l_1 + l_2 + \cdots + l_{s-1} + k_1 + k_2 + \cdots + k_{s-1}} \\
& \times (b_{s-1} c_{s-1})^{-n_{s-1}} \cdots (b_1 c_1)^{-n_1} (b'_{s-1} c'_{s-1})^{-l_{s-1}} \cdots (b'_1 c'_1)^{-l_1} \\
& \times (b'_{s-1} c'_{s-1})^{-k_{s-1}} \cdots (b'_1 c'_1)^{-k_1} \beta_{(n_1, l_1, k_1)}. \tag{5.11}
\end{aligned}$$

Proof :

Equation (5.11) is the result of a s -fold iteration of SBTL with the choice $b = b_i, c = c_i, b' = b'_i, c' = c'_i, b'' = b''_i, c'' = c''_i$ at the i th step. $\square$

Theorem 5.3

If $(\alpha_m, \beta_{(n,l,k)})$ is a SBTP (i.e. related by (5.1)), then

$$\begin{aligned}
& \sum_{\substack{n_s \geq \dots \geq n_1 \geq 0 \\ l_s \geq \dots \geq l_1 \geq 0 \\ k_s \geq \dots \geq k_1 \geq 0}} \frac{q^{n_1^2 + n_2^2 + \dots + n_s^2 + l_1^2 + l_2^2 + \dots + l_s^2 + k_1^2 + k_2^2 + \dots + k_s^2}}{(q; q)_{n_s - n_{s-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_s - l_{s-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{a^{n_1 + n_2 + \dots + n_s + l_1 + l_2 + \dots + l_s + k_1 + k_2 + \dots + k_s} \beta_{(n_1, l_1, k_1)}}{(q; q)_{k_s - k_{s-1}} \cdots (q; q)_{k_2 - k_1}} \\
& = \frac{1}{(aq; q)_\infty^3} \sum_{m=0}^{\infty} q^{3sm^2} a^{3sm} \alpha_m \quad (5.12)
\end{aligned}$$

Proof :

Letting $N, L, K, b_1, \dots, b_s, c_1, \dots, c_s, b'_1, \dots, b'_s, c'_1, \dots, c'_s, b''_1, \dots, b''_s, c''_1, \dots, c''_s$ all tend to infinity in Theorem 5.2, the proof of Theorem 5.3 follows. $\square$

Theorem 5.3 may now be used to embed the triple series Rogers-Ramanujan type identities (5.9) and (5.10) in an infinite family of such identities:

$$\begin{aligned}
& \sum_{\substack{n_s \geq \dots \geq n_1 \geq 0 \\ l_s \geq \dots \geq l_1 \geq 0 \\ k_s \geq \dots \geq k_1 \geq 0}} \frac{q^{n_1^2 + \dots + n_s^2 + l_1^2 + \dots + l_s^2 + k_1^2 + \dots + k_s^2}}{(q; q)_{n_s - n_{s-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_s - l_{s-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{(q; q)_{n_1 + l_1 + k_1}}{(q; q)_{k_s - k_{s-1}} \cdots (q; q)_{k_2 - k_1} (q; q)_{n_1 + l_1} (q; q)_{n_1 + k_1} (q; q)_{l_1 + k_1}} \\
& \times \frac{1}{(q; q)_{n_1} (q; q)_{l_1} (q; q)_{k_1}} = \frac{1}{(q; q)_\infty^2} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm(3s+1) \pmod{6s+3}}}^{\infty} (1 - q^m)^{-1} \quad (5.13)
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{\substack{n_s \geq \dots \geq n_1 \geq 0 \\ l_s \geq \dots \geq l_1 \geq 0 \\ k_s \geq \dots \geq k_1 \geq 0}} \frac{q^{n_1^2 + \dots + n_s^2 + l_1^2 + \dots + l_s^2 + k_1^2 + \dots + k_s^2}}{(q; q)_{n_s - n_{s-1}} \cdots (q; q)_{n_2 - n_1} (q; q)_{l_s - l_{s-1}} \cdots (q; q)_{l_2 - l_1}} \\
& \times \frac{q^{n_1 + \dots + n_s + l_1 + \dots + l_s + k_1 + \dots + k_s}}{(q; q)_{k_s - k_{s-1}} \cdots (q; q)_{k_2 - k_1} (q^2; q)_{n_1 + l_1} (q^2; q)_{n_1 + k_1} (q^2; q)_{l_1 + k_1}} \\
& \times \frac{(q^2; q)_{n_1 + l_1 + k_1}}{(q^2; q)_{n_1} (q^2; q)_{l_1} (q^2; q)_{k_1}} = \frac{1}{(q^2; q)_\infty^2} \prod_{\substack{m=1 \\ m \not\equiv 0, \pm 1 \pmod{6s+3}}}^{\infty} (1 - q^m)^{-1}. \quad (5.14)
\end{aligned}$$

References

- [1] R. P. Agarwal; *Resonance of Ramanujan's Mathematics, Vol. I*, New Age International Publishers, New Delhi (1996).
- [2] A. K. Agarwal, G. E. Andrews and D. M. Bressoud; *The Bailey lattice*, J. Indian Math. Soc.(N.S.), 51 (1987), 57-73.
- [3] G. E. Andrews; *A general theory of identities of Rogers-Ramanujan type*, Bul. Amer. Math. Soc., 80 (1974), 1033-1052.
- [4] G. E. Andrews; *An analytic generalization of the Rogers-Ramanujan identities for odd moduli*, Proc. Nat. Acad. Sci. USA 71 (1974), 4082-4085.
- [5] G. E. Andrews; *Problems and prospects for basic hypergeometric functions*, Theory and Application of Special Functions (Proc. Advanced Sem., Math. Res. Center, Univ. Wisconsin, Madison, Wis., 1975) (R. Askey, ed.), Math. Res. Center, Univ. Wisconsin, Publ. no. 35, Academic Press, New York, 1975, pp. 191-224.
- [6] G. E. Andrews; *The Theory of Partitions*, Encyclopedia of Mathematics and Its Applications, vol. 2, Addison-Wesley Publishing, Massachusetts, 1976.
- [7] G. E. Andrews; *Partition and Durfee dissection*, Amer. J. Math. 101 (1979), no. 3, 735-742.
- [8] G. E. Andrews; *The hard hexagon model and Rogers-Ramanujan identities*, Proc. Natl. Acad. Sci. USA 78 (1981), no. 9, part 1, 5290-5292.
- [9] G. E. Andrews; *Multiple series Rogers-Ramanujan type identities*, Pacific J. Math. 114 (1984), no. 2, 267-283.

- [10] G. E. Andrews; *q-Series : Their Development and Application in Analysis, Number Theory, Combinatorics, Physics and Computer Algebra*, CBMS Regional Conference Series in Mathematics, vol. 66, the American Mathematical Society, Rhode Island, 1986.
- [11] G. E. ANDREWS; *Bailey chains and generalized Lambert series: I. Four identities of Ramanujan*, Illinois J. Math. 36 (1992), 251-274.
- [12] G. E. ANDREWS; *Umbral calculus, Bailey chains and pentagonal number theorems*, J. Combin. Th. (A), 91 (2000), 464-475.
- [13] G. E. ANDREWS; *Baileys transform, lemma, chain and tree*, Special Functions 2000, eds. J. Bustoz et al., Kluwer, (2001), 1-22.
- [14] G. E. ANDREWS; *A Bailey lemma from the quintuple product*, In Number Theory and Discrete Mathematics, A. Agarwal et al. eds, Hindustan Book Agency, (2002), 37-46.
- [15] G. E. Andrews; *The Bailey transform and false theta functions* (submitted).
- [16] G. E. ANDREWS, R. J. BAXTER AND P. J. FORRESTER; *Eight-vertex SOS model and generalized RogersRamanujan-type identities*, J. Stat. Phys. 35 (1984), 193-266.
- [17] G. E. ANDREWS AND A. BERKOVICH; *The WP-Bailey tree and its implications*, J. London Math. Soc., 66 (2002), no. 3, 529-549.
- [18] G. E. ANDREWS AND A. BERKOVICH; *A trinomial analogue of Baileys lemma and $N = 2$ superconformal invariance*, Commun. Math. Phys. 192 (1998), 245-260.
- [19] G. E. ANDREWS AND D. BOWMAN; *The Bailey transform and D.B. Sears*, Quaestiones Mathematicae, 22 (1999), no. 1, 19-26.
- [20] G. E. ANDREWS, A. SCHILLING AND S. O. WARNAAR; *An A2 Bailey lemma and Rogers Ramanujan-type identities*, J. Amer. Math. Soc. 12 (1999), 677-702.
- [21] A. BERKOVICH, B. M. MCCOY AND A. SCHILLING; *$N = 2$ Supersymmetry and Bailey Pairs*, Physica (A) 228 (1996), 33-62.
- [22] A. BERKOVICH, B.M. MCCOY AND P.A. PEARCE; *The perturbations $\phi_{2,1}$ and $\phi_{1,5}$ of the minimal models $M(p, p')$ and the trinomial analogue of Bailey's lemma*, Nucl.Phys.(B), 519 (1998), 597-625.
- [23] W. N. Bailey; *Some identities in combinatorial analysis*, Proc. London Math. Soc. (2) 49 (1947), 421-425.
- [24] W. N. Bailey; *Identities of Rogers-Ramanujan type*, Proc. London Math. Soc. (2) 50 (1948), 1-10.

- [25] D. M. Bressoud; *Some identities for terminating q -series*, Math. Proc. Cambridge Philos. Soc. 89 (1981), no. 2, 211-223.
- [26] D. M. Bressoud; *Change of base in Bailey pairs*, Ramanujan J. 4 (2000), no. 4, 435-453.
- [27] J. L. Burchnall and T. W. Chaundy; *Expansions of Appell's double hypergeometric functions*, Quat. J. of Maths. (Oxford) 11 (1940), 249-270.
- [28] H. Coskum; *An Elliptic BC_n Bailey Lemma and Rogers-Ramanujan Identities Associated To Root Systems*, math.CO/0605653.
- [29] R. Y. Denis, S. N. Singh and S. P. Singh; *On certain transformation formulae for abnormal q -series*, South East Asian J. Math. Math. Sci. 1 (2003), no. 3, 7-19.
- [30] A. Erdélyi; *Higher Transcendental Functions, Vol. I, II*, McGraw-Hill, New York, 1953.
- [31] H. Exton; *q -Hypergeometric Functions and Applications*, Ellis Horwood Series: Mathematics and Its Applications, Ellis Horwood, Chichester, 1983.
- [32] O. FODA AND Y.-H. QUANO; *Virasoro Character Identities from the Andrews Bailey Construction*, Int. J. of Mod. Phys. (A), 12 (1997), 1651-1675.
- [33] G. GASPER AND M. RAHMAN; *Basic Hypergeometric Series*, 2nd ed., Encyclopedia of Mathematics and its Applications, Cambridge Univ. Press, 96 (2004).
- [34] V. K. Jain; *A Study of Certain Hypergeometric Identities*, Ph.D. thesis, University of Roorkee, Roorkee.
- [35] V. K. Jain; *Some expansions involving basic hypergeometric functions of two variables*, Pacific J. Math. 91 (1980), no. 2, 349-361.
- [36] V. K. Jain; *Certain transformations of basic hypergeometric series and their applications*, Pacific J. Math. 101 (1982), no. 2, 333-349.
- [37] C. M. JOSHI AND Y. VYAS; *Extensons of Bailey's transform and applications*, Int. J. Math. Math. Sci. 2005 (2005), no. 12, 1909-1923.
- [38] J. LOVEJOY; *A Bailey Lattice*, Proc. Amer. Math. Soc. 132 (2004), 1507-1516.
- [39] P. A. MACMOHON; *Combinatory Analysis, Vol. II*, Cambridge Univ. Press, London, (1918).
- [40] E. B. McBride; *Obtaining Generating Functions*, Springer Tracts in Natural Philosophy, vol. 21, Springer, New York, 1971.
- [41] P. PAULE; *On identities of the RogersRamanujan type*, J. Math. Anal. Appl. 107 (1985), 255-284.

- [42] P. PAULE; *A note on Baileys lemma*, J. Combin. Theory Ser. (A) 44 (1987), 164-167.
- [43] P. PAULE; *The concept of Bailey chains*, Publ. I.R.M.A. Strasbourg, 358/S-18 (1988), 53-76.
- [44] E. D. Rainville; *Special Functions*, MacMillan Co., New York, 1967.
- [45] S. RAMANUJAN; *Ramanujan's Lost Notebook, Vol. I*, George Andrews and Bruce Berndt, editors, Springer, (2005).
- [46] L. J. ROGERS; *Second memoirs on the expansion of certain infinite products*, Proc. London Math. Soc., 25 (1894), 318-343.
- [47] L. J. ROGERS; *On two theorems of combinatory analysis and some allied identities*, Proc. London Math. Soc., 16 (1917), 315-336.
- [48] O. Shankar; *A Watson's theorem for the double hypergeometric series*, Ganita, 27 (1976), 73-78.
- [49] A. SCHILLING AND L. DEKA; *Non-unitary minimal models, Bailey's lemma and $N = 1, 2$ superconformal algebras*, Commun. Math. Phys. 260 (2005), 711-725.
- [50] A. SCHILLING AND S. O. WARNAAR; *A higher level Bailey lemma: proof and application*, The Ramanujan Journal 2 (1998), 327-349.
- [51] A. SCHILLING AND S. O. WARNAAR; *Inhomogeneous lattice paths, generalized Kostka polynomials and A_{n-1} supernomials*, Commun. Math. Phys. 202 (1999), 359-401.
- [52] A. SCHILLING AND S. O. WARNAAR; *Conjugate Bailey pairs. From configuration sums and fractional-level string functions to Bailey's lemma*, Contemp. Math. 297 (2002), 227-255.
- [53] A. SCHILLING, A. BERKOVICH, B. M. MCCOY AND S. O. WARNAAR; *Bailey flows and Bose-Fermi identities for the conformal coset models $(A_1^{(1)})_N \times (A_1^{(1)})_{N'}/(A_1^{(1)})_{N+N'}$* , Nucl. Phys. B499 [PM] (1997), 621-649.
- [54] A. SCHILLING AND S. O. WARNAAR; *A Higher-level Bailey Lemma*, Int. J. of Mod. Phys. (B) 11 (1997), 189-195.
- [55] A.V. SILLS; *On identities of Rogers-Ramanujan type*, Ramanujan J., to appear.
- [56] A.V. SILLS; *q -difference equations and identities of the Rogers-Ramanujan-Bailey type*, J. Diff. Equ. Appl. 10 (2004), 1069-1084.
- [57] A.V. SILLS; *Identities of the Rogers-Ramanujan-Bailey type*, J. Math. Anal. Appl. 308/2 (2005), 669-688.

- [58] A.V. SILLS; *On identities of Rogers-Ramanujan-Slater type*, Int. J. Num. Th., to appear.
- [59] S. P. Singh; *Certain transformation formulae for q -series*, Indian J. Pure Appl. Math. 31 (2000), no. 10, 1369-1377.
- [60] U. B. Singh; *A note on a transformation of Bailey*, Quart. J. Math. Oxford Ser. (2) 45 (1994), no. 177, 111-116.
- [61] L. J. Slater; *A new proof of Rogers transformations of infinite series*, Proc. London Math. Soc. (2) 53 (1951), 460-475.
- [62] L. J. Slater; *Further identities of Rogers-Ramanujan type*, Proc. London Math. Soc. (2) 54 (1952), 147-167.
- [63] L. J. Slater; *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [64] V. P. SPIRIDONOV ; *An elliptic incarnation of the Bailey chain*, Int. Math. Res. Not., 37 (2002), 1945-1977.
- [65] H. M. Srivastava and H. L. Manocha; *A Treatise on Generating Functions*, Ellis Horwood Series: Mathematics and Its Applications, Ellis Horwood, Chichester, 1984.
- [66] A. Verma and V. K. Jain; *Transformations between basic hypergeometric series on different base and identities of Rogers-Ramanujan type*, J. Math. Anal. Appl. 76 (1980), no. 1, 230-269.
- [67] A. Verma and V. K. Jain; *Transformations of non-terminating basic hypergeometric series, their contour integrals and applications to Rogers-Ramanujan identities*, J. Math. Anal. Appl. 87 (1982), no. 1, 9-44.
- [68] S. O. WARNAAR; *Fermionic solution of the Andrews-Baxter-Forrester model. II. Proof of Melzer's polynomial identities*, J. Stat. Phys., 84 (1996), 49-83.
- [69] S.O. WARNAAR; *A note on the trinomial analogue of Bailey's lemma*, J. Combin. Th. (A), 81 (1998), 114-118.
- [70] S.O. WARNAAR; *Supernomial coefficients, Bailey's lemma and Rogers-Ramanujan-type identities*, A survey of results and open problems Séminaire Lotharingien de Combinatoire 42 (1999), Art. B42n 22p.
- [71] S.O. WARNAAR; *50 Years of Bailey's lemma*, in Algebraic Combinatorics and Applications, A. Betten et al. eds., Springer, Berlin, (2001), 333-347.
- [72] S.O. WARNAAR; *Extensions of the well-poised and elliptic well-poised Bailey lemma*, Indagationes Mathematicae New Series 14 (2003), 571-588.

- [73] S. O. WARNAAR; *Partial theta functions. I. Beyond the lost notebook*, Proc. London Math. Soc. 87 (2003), 363-395.
- [74] S.O. WARNAAR; *The Bailey lemma and Kostka polynomials*, J. Alg. Combin., 20 (2004), 131-171.

106, Arihant Nagar, Kalka Mata Road,
Udaipur-313 001, Rajasthan, INDIA.

— —

Department of Mathematics and Statistics,
University College of Science,
Mohanlal Sukhadia University,
Udaipur-313 001, Rajasthan, INDIA.
E-Mail : *yashoverdhan@yahoo.com*